arXiv:0712.2567v1 [cs.DM] 17 Dec 2007

Mathematical Problems of Computer Science 23, 2004, 127–129.

On Lower Bound for $W(K_{2n})$

Rafael R. Kamalian and Petros A. Petrosyan
Institute for Informatics and Automation Problems of NAS of RA
e-mails rrkamalian@yahoo.com, pet_petros@yahoo.com

Abstract

The lower bound $W(K_{2n}) \geq 3n-2$ is proved for the greatest possible number of colors in an interval edge coloring of the complete graph K_{2n}.

Let $G = (V(G), E(G))$ be an undirected graph without loops and multiple edges [1], $V(G)$ and $E(G)$ be the sets of vertices and edges of G, respectively. The degree of a vertex $x \in V(G)$ is denoted by $d_G(x)$, the maximum degree of a vertex of G-by $\Delta(G)$, and the chromatic index of G-by $\chi'(G)$. A graph is regular, if all its vertices have the one degree. If α is a proper edge coloring of the graph G, then $\alpha(e)$ denotes the color of an edge $e \in E(G)$ in the coloring α.

Let α be a proper coloring of edges of G with colors $1, 2, \ldots, t$. α is interval [2], if for each color $i, 1 \leq i \leq t$, there exists at least one edge $e_i \in E(G)$ with $\alpha(e_i) = i$ and the edges incident with each vertex $x \in V(G)$ are colored by $d_G(x)$ consecutive colors.

A graph G is interval-colorable if there is $t \geq 1$ for which G has an interval edge coloring α with colors $1, 2, \ldots, t$. The set of all interval-colorable graphs is denoted by $\mathcal{N}$ [2].

For $G \in \mathcal{N}$ we denote by $W(G)$ the greatest value of t, for which G has an interval edge coloring α with colors $1, 2, \ldots, t$.

It was proved [3] for bipartite graphs that verification whether $G \in \mathcal{N}$ is NP-complete [4,5].

It was proved [2] that if G has no triangle and $G \in \mathcal{N}$ then $W(G) \leq |V(G)| - 1$. It follows from here that if G is bipartite and $G \in \mathcal{N}$ then $W(G) \leq |V(G)| - 1$ [2].

For graphs which can contain a triangle the following results hold:

Theorem 1 [6]. If $G \in \mathcal{N}$ is a graph with nonempty set of edges then $W(G) \leq 2\,|V(G)| - 3$.

Theorem 2 [7]. If $G \in \mathcal{N}$ and $|V(G)| \geq 3$ then $W(G) \leq 2\,|V(G)| - 4$.

Non-defined conceptions and notations can be found in [1,6,8,9].

In [2] there was proved the following

Theorem 3. If $G \in \mathcal{N}$ then $\chi'(G) = \Delta(G)$.

Corollary 1 [2]. For a regular graph G $G \in \mathcal{N}$ iff $\chi'(G) = \Delta(G)$.

A well-known inequality $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$ for the chromatic index of an arbitrary graph G was proved in [10]. It was proved [11] for regular graphs that verification whether $\chi'(G) = \Delta(G)$ is NP-complete.Therefore we can conclude from the **Corollary 1** that the problem

"Whether a given regular graph belongs to the set $\mathcal{N}$ or not?" is also NP-complete.

From the results of [10] and **Corollary 1** it follows that for any odd p $K_p \notin \mathcal{N}$. It is not difficult to see that $\chi'(K_{2n}) = \Delta(K_{2n})$ (it can be obtained from the theorem 9.1 of [1]). Consequently from the **Corollary 1** we obtain

Theorem 4. For any $n \in N \quad K_{2n} \in \mathcal{N}$.

It is not difficult, using the results of [6], to obtain the following

Theorem 5. For any $n \in N \quad W(K_{2n}) \geq 2n - 1 + \lfloor \log_2 (2n-1) \rfloor$.

The main result of this paper consists of the following

Theorem 6. For any $n \in N \quad W(K_{2n}) \geq 3n - 2$.

Proof. Obviously, for $n \leq 3$ the estimate of the **Theorem 6** coincides with that one of the **Theorem 5**.

Now assume that $n \geq 4$.

Consider a graph K_{2n} with $V(K_{2n}) = \{u_1, u_2, ..., u_{2n}\}$ and
$E(K_{2n}) = \{(u_i, u_j) \mid \ u_i \in V(K_{2n}),\ u_j \in V(K_{2n}),\ i < j\}$.

Define a coloring α of the edges of the graph K_{2n} in the following way:

for $i = 1, ..., \left\lfloor \frac{n}{2} \right\rfloor, j = 2, ..., n, i < j, i + j \leq n + 1$

$$\alpha\left((u_i, u_j)\right) = i + j - 2;$$

for $i = 2, ..., n-1, j = \left\lfloor \frac{n}{2} \right\rfloor + 2, ..., n, i < j, i + j \geq n + 2$

$$\alpha\left((u_i, u_j)\right) = i + j + n - 3;$$

for $i = 3, ..., n, j = n + 1, ..., 2n - 2, j - i \leq n - 2$

$$\alpha\left((u_i, u_j)\right) = n + j - i;$$

for $i = 1, ..., n, j = n + 1, ..., 2n, j - i \geq n$

$$\alpha\left((u_i, u_j)\right) = j - i;$$

for $i = 2, ..., 1 + \left\lfloor \frac{n-1}{2} \right\rfloor, j = n + 1, ..., n + \left\lfloor \frac{n-1}{2} \right\rfloor,\ j - i = n - 1$

$$\alpha\left((u_i, u_j)\right) = 2(i - 1);$$

for $i = \left\lfloor \frac{n-1}{2} \right\rfloor + 2, ..., n,\ j = n + 1 + \left\lfloor \frac{n-1}{2} \right\rfloor, ..., 2n - 1,\ j - i = n - 1$

$$\alpha\left((u_i, u_j)\right) = i + j - 2;$$

for $i = n + 1, ..., n + \left\lfloor \frac{n}{2} \right\rfloor - 1, j = n + 2, ..., 2n - 2, i < j, i + j \leq 3n - 1$

$$\alpha\left((u_i, u_j)\right) = i + j - 2n;$$

for $i = n + 1, ..., 2n - 1, j = n + \left\lfloor \frac{n}{2} \right\rfloor + 1, ..., 2n, i < j, i + j \geq 3n$

$$\alpha\left((u_i, u_j)\right) = i + j - n - 1.$$

It is not difficult to see that α is an interval edge coloring of the graph K_{2n} with colors $1, 2, ..., 3n - 2$.

The proof is complete.

References

[1] F. Harary, Graph Theory, Addison-Wesley, Reading, MA,1969.

[2] A.S. Asratian, R.R. Kamalian, Interval colorings of edges of a multigraph, *Appl. Math.*5 , 25-34,1987.

[3] S.V. Sevastianov, On interval colourability of edges of a bipartite graph, *Meth. Of Discr. Anal.* In solution of external problems. The Institute of Mathematics of the Siberian Branch of the Academy of Sciences of USSR. Novosibirsk, N50, 61-72, 1990.

[4] S. Cook, The complexity of theorem-proving procedures. In *Proc.3rd ACM Symp.* on Theory of Computing, 151-158, 1971.

[5] R.M. Karp, Reducibility among Combinatorial Problems, in “Complexity of Computer Computations” (R.E. Miller and J.W. Thatcher, Eds.), pp. 85-103, New York, Plenum, 1972.

[6] R.R. Kamalian, Interval Edge Colorings of Graphs, Doctoral dissertation, Novosibirsk, 1990.

[7] K. Giaro, M. Kubale, M. Malafiejski, Consecutive colorings of the edges of general graphs, *Discr. Math.* 236 , 131-143,2001.

[8] A.A. Zykov, Theory of finite graphs, Novosibirsk, Nauka, 1969.

[9] A.S. Asratian, R.R. Kamalian, Investigation on interval edge colorings of graphs, *J. Combin. Theory Ser.* B 62, 34-43,1994.

[10] V.G. Vizing, The chromatic index of a multigraph, *Kibernetika* 3, 29-39, 1965.

[11] I. Holyer, The NP-completeness of edge coloring, *SIAM J. Comput.* 10, N4, 718-720, 1981.